

DEPARTMENT OF PUBLIC HEALTH AND ENVIRONMENT

Water Quality Control Commission

REGULATION NO. 34 - CLASSIFICATIONS AND NUMERIC STANDARDS FOR SAN JUAN RIVER AND DOLORES RIVER BASINS

5 CCR 1002-34

[Editor's Notes follow the text of the rules at the end of this CCR Document.]

34.6 TABLES

(3) Table Value Standards

In certain instances in the tables in Appendix 34-1, the designation "TVS" is used to indicate that for a particular parameter a "table value standard" has been adopted. This designation refers to numerical criteria set forth in the Basic Standards and Methodologies for Surface Water. The criteria for which the TVS are applicable are on the following table.

**TABLE VALUE STANDARDS
(Concentrations in µg/l unless noted)**

| PARAMETER ⁽¹⁾ | TABLE VALUE STANDARDS ⁽²⁾⁽³⁾ |
|--------------------------|--|
| Aluminum (T) | Acute = $e^{(1.3695[\ln(\text{hardness})]+1.8308)}$ pH equal to or greater than 7.0 Chronic = $e^{(1.3695[\ln(\text{hardness})]-0.1158)}$ pH less than 7.0 Chronic = $e^{(1.3695[\ln(\text{hardness})]-0.1158)}$ or 87, whichever is less |
| Ammonia ⁽⁴⁾ | <p>Cold Water = (mg/l as N) Total</p> $acute = \frac{0.275}{1 + 10^{7.204 - pH}} + \frac{39.0}{1 + 10^{pH - 7.204}}$ $chronic = \left(\frac{0.0577}{1 + 10^{7.688 - pH}} + \frac{2.487}{1 + 10^{pH - 7.688}} \right) * MIN(2.85, 1.45 * 10^{0.028(25 - T)})$ <p>Warm Water = (mg/l as N) Total</p> $acute = \frac{0.411}{1 + 10^{7.204 - pH}} + \frac{58.4}{1 + 10^{pH - 7.204}}$ $chronic (Apr1 - Aug31) = \left(\frac{0.0577}{1 + 10^{7.688 - pH}} + \frac{2.487}{1 + 10^{pH - 7.688}} \right) * MIN(2.85, 1.45 * 10^{0.028(25 - T)})$ $chronic (Sep1 - Mar31) = \left(\frac{0.0577}{1 + 10^{7.688 - pH}} + \frac{2.487}{1 + 10^{pH - 7.688}} \right) * 1.45 * 10^{0.028 * (25 - MAX(T, 7))}$ |

| | | | | | | |
|------------------------------|---|--|---|-------------------|---------------------------|------|
| Cadmium | $\text{Acute(warm)}^{(5)} = (1.136672 - (\ln(\text{hardness}) * 0.041838)) * e^{(0.9789 * \ln(\text{hardness}) - 3.443)}$ $\text{Acute(cold)}^{(5)} = (1.136672 - (\ln(\text{hardness}) * 0.041838)) * e^{(0.9789 * \ln(\text{hardness}) - 3.866)}$ $\text{Chronic} = (1.101672 - (\ln(\text{hardness}) * 0.041838)) * e^{(0.7977 * \ln(\text{hardness}) - 3.909)}$ $\text{Acute} = (1.136672 - [\ln(\text{hardness}) * (0.041838)]) * e^{(0.9151 * \ln(\text{hardness}) - 3.1485)}$ $\text{Acute(Trout)} = (1.136672 - [\ln(\text{hardness}) * (0.041838)]) * e^{(0.9151 * \ln(\text{hardness}) - 3.6236)}$ $\text{Chronic} = (1.101672 - [\ln(\text{hardness}) * (0.041838)]) * e^{(0.7998 * \ln(\text{hardness}) - 4.4451)}$ | | | | | |
| Chromium III ⁽⁵⁶⁾ | Acute = $e^{(0.819[\ln(\text{hardness})] + 2.5736)}$ Chronic = $e^{(0.819[\ln(\text{hardness})] + 0.5340)}$ | | | | | |
| Chromium VI ⁽⁵⁶⁾ | Acute = 16 Chronic = 11 | | | | | |
| Copper | Acute = $e^{(0.9422[\ln(\text{hardness})] - 1.7408)}$ Chronic = $e^{(0.8545[\ln(\text{hardness})] - 1.7428)}$ | | | | | |
| Lead | Acute = $(1.46203 - [(\ln(\text{hardness}) * (0.145712))] * e^{(1.273[\ln(\text{hardness})] - 1.46)}$ Chronic = $(1.46203 - [(\ln(\text{hardness}) * (0.145712))] * e^{(1.273[\ln(\text{hardness})] - 4.705)}$ | | | | | |
| Manganese | Acute = $e^{(0.3331[\ln(\text{hardness})] + 6.4676)}$ Chronic = $e^{(0.3331[\ln(\text{hardness})] + 5.8743)}$ | | | | | |
| Nickel | Acute = $e^{(0.846[\ln(\text{hardness})] + 2.253)}$ Chronic = $e^{(0.846[\ln(\text{hardness})] + 0.0554)}$ | | | | | |
| Selenium ⁽⁶⁷⁾ | Acute = 18.4 Chronic = 4.6 | | | | | |
| Silver | Acute = $\frac{1}{2}e^{(1.72[\ln(\text{hardness})] - 6.52)}$ Chronic = $e^{(1.72[\ln(\text{hardness})] - 9.06)}$ Chronic(Trout) = $e^{(1.72[\ln(\text{hardness})] - 10.51)}$ | | | | | |
| Temperature | TEMPERATURE TIER | TIER CODE | SPECIES EXPECTED TO BE PRESENT | APPLICABLE MONTHS | TEMPERATURE STANDARD (°C) | |
| | | | | | MWAT | DM |
| | Cold Stream Tier 1 | CS-I | brook trout, cutthroat trout | June – Sept. | 17.0 | 21.7 |
| | | | | Oct. – May | 9.0 | 13.0 |
| | Cold Stream Tier 2 | CS-II | all other cold-water species | April – Oct. | 18.3 | 24.3 |
| | | | | Nov. – March | 9.0 | 13.0 |
| | Cold Lakes | CL | brook trout, brown trout, cutthroat trout, lake trout, rainbow trout, Arctic grayling, sockeye salmon | April – Dec. | 17.0 | 21.2 |
| | | | | Jan. – March | 9.0 | 13.0 |
| | Cold Large Lakes (>100 acres surface area) | CLL | rainbow trout, brown trout, lake trout | April – Dec. | 18.3 | 24.2 |
| | | | | Jan. – March | 9.0 | 13.0 |
| Warm Stream Tier 2 | WS-II | brook stickleback, central stoneroller, creek chub, longnose dace, northern redbelly dace, finescale dace, razorback sucker, white sucker, mountain sucker | March – Nov. | 27.5 | 28.6 | |
| | | | Dec. – Feb. | 13.8 | 25.2 | |
| Warm Stream Tier 3 | WS-III | all other warm-water species | March – Nov. | 28.7 | 31.8 | |
| | | | Dec. – Feb. | 14.3 | 24.9 | |

| | | | | | | |
|---------|---|----|--|--------------|------|------|
| | Warm Lakes | WL | black crappie, bluegill, common carp, gizzard shad, golden shiner, largemouth bass, northern pike, pumpkinseed, sauger, smallmouth bass, spottail shiner, stonecat, striped bass, tiger muskellunge, walleye, wiper, white bass, white crappie, yellow perch | April – Dec. | 26.2 | 29.3 |
| | | | | Jan. – March | 13.1 | 24.1 |
| Uranium | Acute = $e^{(1.1021[\ln(\text{hardness})]+2.7088)}$ Chronic = $e^{(1.1021[\ln(\text{hardness})]+2.2382)}$ | | | | | |
| Zinc | Acute = $0.978 * e^{(0.9094[\ln(\text{hardness})]+0.9095)}$ Chronic = $0.986 * e^{(0.9094[\ln(\text{hardness})]+0.6235)}$ Where hardness is less than 102 mg/L CaCO ³ and mottled sculpin are expected to be present: Chronic (sculpin) = $e^{(2.140[\ln(\text{hardness})]-5.084)}$ | | | | | |

TABLE VALUE STANDARDS - FOOTNOTES

- (1) Metals are stated as dissolved unless otherwise specified.
- (2) Hardness values to be used in equations are in mg/L as calcium carbonate and shall be no greater than 400 mg/L, except for aluminum for which hardness shall be no greater than 220 mg/L. The hardness values used in calculating the appropriate metal standard should be based on the lower 95 per cent confidence limit of the mean hardness value at the periodic low flow criteria as determined from a regression analysis of site-specific data. Where insufficient site-specific data exists to define the mean hardness value at the periodic low flow criteria, representative regional data shall be used to perform the regression analysis. Where a regression analysis is not appropriate, a site-specific method should be used. In calculating a hardness value, regression analyses should not be extrapolated past the point that data exist.
- (3) Both acute and chronic numbers adopted as stream standards are levels not to be exceeded more than once every three years on the average.
- (4) For acute conditions the default assumption is that salmonids could be present in cold water segments and should be protected, and that salmonids do not need to be protected in warm water segments. For chronic conditions, the default assumptions are that early life stages could be present all year in cold water segments and should be protected. In warm water segments the default assumption is that early life stages are present and should be protected only from April 1 through August 31. These assumptions can be modified by the commission on a site-specific basis where appropriate evidence is submitted.
- (5) The acute(warm) cadmium equation applies to segments classified as Aquatic Life Warm Class 1 or 2. The acute(cold) cadmium equation applies to segments classified as Aquatic Life Cold Class 1 or 2.
- (56) Unless the stability of the chromium valence state in receiving waters can be clearly demonstrated, the standard for chromium should be in terms of chromium VI. In no case can the sum of the instream levels of Hexavalent and Trivalent Chromium exceed the

water supply standard of 50 µg/l total chromium in those waters classified for domestic water use.

- (67) Selenium is a bioaccumulative metal and subject to a range of toxicity values depending upon numerous site-specific variables.

34.51 STATEMENT OF BASIS, SPECIFIC STATUTORY AUTHORITY AND PURPOSE; DECEMBER 9, 2019 RULEMAKING; FINAL ACTION JANUARY 13, 2020; EFFECTIVE DATE JUNE 30, 2020

The provisions of C.R.S. 25-8-202(1)(a), (b) and (2); 25-8-203; 25-8-204; and 25-8-402; provide the specific statutory authority for adoption of these regulatory amendments. The commission also adopted in compliance with 24-4-103(4) C.R.S. the following statement of basis and purpose.

BASIS AND PURPOSE

A. Aquatic Life Standards for Cadmium

Cadmium is a naturally-occurring element frequently found alongside other metals, and numerous treatment techniques are available to remove cadmium from wastewater. Cadmium has both acute and chronic effects on aquatic life, and can negatively impact survival, growth, reproduction, immune and endocrine systems, development, and behavior.

The commission revised the hardness-based cadmium table value standards to protect the Aquatic Life use. The updated standards incorporate toxicity data that have become available since the cadmium standards were last updated in the 2005 Regulation No. 31 rulemaking hearing. The updated standards are based on the United States Environmental Protection Agency's (EPA) "Aquatic Life Ambient Water Quality Criteria – 2016" and toxicity data that have become available since EPA's recommended criteria were released in 2016.

The updated standards include two acute equations (acute(cold) and acute(warm)) and one chronic equation. The acute(cold) and chronic equations are the same as the acute and chronic criteria recommended by EPA in 2016. The acute(cold) equation, which is lowered to protect trout, is protective of trout and other sensitive cold water species and applies in segments classified as Aquatic Life Cold Class 1 or 2. The acute(warm) equation, which is not lowered to protect trout, is protective of warm water species and applies in segments classified as Aquatic Life Warm Class 1 or 2. The chronic equation is protective of both cold and warm water aquatic life and applies in segments classified as either Aquatic Life Cold Class 1 or 2 or Aquatic Life Warm Class 1 or 2.

Compared to the previous cadmium table value standards, the updated standards are generally less stringent. The acute(cold) standard is less stringent than the previous acute(trout) standard when water hardness is greater than 45 mg/L CaCO₃. The acute(warm) equation is less stringent than the previous acute standard when water hardness is greater than 101 mg/L CaCO₃. The updated chronic equation is less stringent than the previous chronic standard at all water hardness values.

In the past, Colorado has had separate acute equations for waters with trout and waters without trout. The updated standards include separate acute equations for cold waters (both with and without trout) and warm waters. This change in approach is due to the addition of toxicity data showing that sculpin, which inhabit cold waters, are also sensitive to cadmium. To ensure protection of sculpin and other sensitive cold water aquatic life in waters where trout are absent, the acute(cold) equation applies to all cold waters. As a result, the acute trout (tr) qualifier for cadmium is no longer needed on select cold water segments and was deleted from all segments where it had applied.

During the 2017 basin review, the commission adopted EPA's 2016 recommended criteria as site-specific standards in select cold water segments. The updated table value standards for cold waters are the same as EPA's 2016 recommended criteria. Therefore, to reflect the commission's state-wide adoption of the updated table value standards, the cadmium "SSE" were replaced with "TVS" on the following segments:

Animas Florida: 3a, 3c, 4a, 4b, 6 and 9

Dolores River: 9

B. Clarifications to Appendix 34-1

To improve the clarity and usability of the tables, an acronym list was added to the front of Appendix 34-1 and the footnote referencing Section 34.6 was also simplified.
